

WELL-SAMPLED FAR-INFRARED SPECTRAL ENERGY DISTRIBUTIONS OF $z \sim 2$ GALAXIES: EVIDENCE FOR SCALED UP COOL GALAXIES

ADAM MUZZIN¹, PIETER VAN DOKKUM¹, MARISKA KRIEK², IVO LABBÉ⁴, IARA CURY¹, DANILO MARCHESINI³, & MARIJN FRANK⁵

Draft version October 18, 2010

ABSTRACT

We present an analysis of the far-infrared (FIR) spectral energy distributions (SEDs) of two massive K-selected galaxies at $z = 2.122$ and $z = 2.024$ detected at $24\mu\text{m}$, $70\mu\text{m}$, $160\mu\text{m}$ by Spitzer, $250\mu\text{m}$, $350\mu\text{m}$, $500\mu\text{m}$ by BLAST, and $870\mu\text{m}$ by APEX. The large wavelength range of these observations and the availability of spectroscopic redshifts allow us to unambiguously identify the peak of the redshifted thermal emission from dust at $\sim 300\mu\text{m}$. The SEDs of both galaxies are reasonably well fit by synthetic templates of local galaxies with $L_{\text{IR}} \sim 10^{11}L_{\odot} - 10^{12}L_{\odot}$ yet both galaxies have $L_{\text{IR}} \sim 10^{13}L_{\odot}$. This suggests that these galaxies are not high redshift analogues of the Hyper-LIRGs/ULIRGs used in local templates, but are instead "scaled up" versions of local ULIRGs/LIRGs. Several lines of evidence point to both galaxies hosting an AGN; however, the relatively cool best fit templates and the optical emission line ratios suggest the AGN is not the dominant source heating the dust. For both galaxies the star formation rate determined from the best-fit FIR SEDs ($\text{SFR}(L_{\text{IR}})$) agrees with the SFR determined from the dust corrected $\text{H}\alpha$ luminosity ($\text{SFR}(\text{H}\alpha)$) to within a factor of ~ 2 ; however, when the SFR of these galaxies is estimated using only the observed $24\mu\text{m}$ flux and the standard luminosity-dependent template method ($\text{SFR}(24\mu\text{m})$), it systematically overestimates the SFR by as much as a factor of 6. A larger sample of 24 K-selected galaxies at $z \sim 2.3$ drawn from the Kriek et al. (2008) GNIRS sample shows the same trend between $\text{SFR}(24\mu\text{m})$ and $\text{SFR}(\text{H}\alpha)$. Using that sample we show that $\text{SFR}(24\mu\text{m})$ and $\text{SFR}(\text{H}\alpha)$ are in better agreement when $\text{SFR}(24\mu\text{m})$ is estimated using the log average of local templates rather than selecting a single luminosity-dependent template, because this incorporates lower luminosity templates. The better agreement between SFRs from lower luminosity templates suggests that the FIR SEDs of the BLAST-detected galaxies may be typical for massive star forming galaxies at $z \sim 2$, and that the majority are scaled up versions of lower luminosity local galaxies.

Subject headings: infrared: galaxies

1. INTRODUCTION

It is well known that in the local universe the obscuration towards star forming regions is correlated with star formation rate (SFR; e.g., Wang & Heckman 1996; Hopkins et al. 2001; Buat et al. 2005, 2007), with the most active galaxies frequently being the most dust obscured. In the most extreme galaxies, the star forming regions can become optically-thick (e.g., Genzel et al. 1998), which makes mid- and far-infrared (MIR, FIR) data critical for a complete metric of their SFR. At $z \sim 2$, where the average massive galaxy is forming stars at rates $\sim 100 - 1000$ times higher than in the local universe (e.g., Pérez-González et al. 2005; Juneau et al. 2005; Damen et al. 2009), MIR- and FIR-determined SFRs are critical for understanding the buildup of today's massive galaxies.

The advent of the MIPS instrument onboard *Spitzer* has facilitated the first deep photometry at $24\mu\text{m}$ and

numerous studies have used these data to infer the star formation history of the universe up to $z \sim 2$ (e.g., Le Floc'h et al. 2005; Pérez-González et al. 2005; Caputi et al. 2007; Reddy et al. 2008; Magnelli et al. 2009). These studies suggest that obscured star formation dominates in massive galaxies at $z \sim 2$; however, the conversion from observed $24\mu\text{m}$ flux to a total infrared luminosity (L_{IR}), and subsequently a SFR at $z \sim 2$ requires significant extrapolation based on local FIR templates.

The most common method for determining the L_{IR} of a galaxy from $24\mu\text{m}$ photometry is to artificially redshift local luminosity-dependent MIR/FIR templates to the redshift of the galaxy and then choose the L_{IR} of the template that predicts a $24\mu\text{m}$ flux closest to the observed $24\mu\text{m}$ flux. By applying this method, or more sophisticated but related versions of it, some authors have found that the local templates produce estimates of L_{IR} and $\text{SFR}(L_{\text{IR}})$ for distant galaxies that are in agreement with other SFR indicators (e.g., Elbaz et al. 2002; Marcellac et al. 2006; Reddy et al. 2006; Daddi et al. 2007a; Papovich et al. 2009; Kartaltepe et al. 2010), while others have argued that this method tends to systematically overestimate the L_{IR} and hence the SFR (e.g., Papovich et al. 2007; Rigby et al. 2008; Murphy et al. 2009). Some of these differences can probably be reconciled by the different selection criteria (e.g., Reddy et al. 2010), redshift range (e.g., Murphy et al. 2009) and luminosities

¹ Department of Astronomy, Yale University, New Haven, CT, 06520-8101; adam.muzzin@yale.edu

² Department of Astrophysical Sciences, Princeton University, Princeton, NJ, 08544

³ Department of Physics and Astronomy, Tufts University, Medford, MA 06520, USA

⁴ Hubble Fellow, Carnegie Observatories, 813 Santa Barbara Street, Pasadena, CA, 91101

⁵ Leiden Observatory, Leiden University, PO Box 9513, 2300 RA Leiden, Netherlands

(e.g., Papovich et al. 2007) of the above samples; however, the reliability of observed $24\mu\text{m}$ fluxes as a metric of the L_{IR} and SFR over the full range of galaxy masses and SFRs at $z \sim 2$ remains unclear.

The next step will be to directly observe the cold dust SED, rather than extrapolate it from the rest-frame $6 - 10\mu\text{m}$ region, which can be complicated by a superposition of Polycyclic Aromatic Hydrocarbon (PAH) emission lines and silicate absorption features. The peak of the cold dust emission in FIR luminous galaxies in the local universe occurs between $40\mu\text{m}$ to $150\mu\text{m}$ which corresponds to $120\mu\text{m}$ to $450\mu\text{m}$ at $z \sim 2$. In the next few years the *Herschel* telescope will provide unprecedented amounts of data at these wavelengths; however, the recent flight of the BLAST telescope (Devlin et al. 2009) has already provided some of the first deep observations of distant galaxies at $250\mu\text{m}$, $350\mu\text{m}$ and $500\mu\text{m}$.

In this paper we take advantage of the BLAST observations of the Extended Chandra Deep Fields South field (ECDFS) to directly measure the FIR SEDs of $z \sim 2$ galaxies. Two star forming galaxies in the Kriek et al. (2008) NIR spectroscopic sample are located in that field and are detected or have upper limits at $24\mu\text{m}$, $70\mu\text{m}$, and $160\mu\text{m}$ in the FIDEL (Magnelli et al. 2009) and SWIRE (Lonsdale et al. 2003) surveys, and are also detected by BLAST (Dye et al. 2009). The ECDFS was recently surveyed by the LABOCA instrument on the APEX telescope, and both galaxies were detected at $870\mu\text{m}$ (Weiss et al. 2009). By combining these data sets we have produced some of the first FIR SEDs that sample the cool dust bump of $z \sim 2$ galaxies with good resolution and allow us to constrain their L_{IR} directly from the FIR SED. Throughout this paper we assume a $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_m = 0.3$, $\Omega_\Lambda = 0.7$ cosmology when computing luminosity distances.

2. GALAXY SAMPLE AND DATA

The NIR spectroscopic sample of Kriek et al. (2008) is a magnitude-limited ($K \leq 19.7$ Vega) sample of 36 galaxies with a median $z \sim 2.3$. The sample is dominated by massive galaxies ($M > 10^{11} M_\odot$), approximately 50% of which have detected emission lines. Two of the galaxies in the sample are extremely bright at $24\mu\text{m}$ ($S_{24\mu\text{m}} > 0.5 \text{ mJy}$), and both are located in the ECDFS (ID#’s ECDFS-4511 and ECDFS-12514 from Kriek et al. 2008). Both galaxies have strong $H\alpha$ emission and Kriek et al. (2008) used this to determine spectroscopic redshifts of $z = 2.122$ and $z = 2.024$. The ID numbers of the galaxies in the BLAST survey of the ECDFS are BLAST J033138-274122 and BLAST J033242-275511. They are also detected at $870\mu\text{m}$ in the 200-hr map from the LABOCA array on the APEX telescope (Weiss et al. 2009, ID#’s LESS J033139.6-274120, and LESS J033243.3-275517).

Using the v0.5 public-release FIDEL⁶ MIPS imaging data of the ECDFS we have measured total $24\mu\text{m}$ and $70\mu\text{m}$ fluxes for these galaxies. For the $24\mu\text{m}$ data the fluxes were measured within $11''$ diameter apertures using the SExtractor package (Bertin & Arnouts 1996) and an aperture correction from the MIPS data handbook⁷ was applied. For the $70\mu\text{m}$ data we used the $24\mu\text{m}$ centroid and measured fluxes within $32''$ apertures and ap-

plied the aperture correction from the MIPS data handbook. The ECDFS was observed at $160\mu\text{m}$ as part of the SWIRE survey (Lonsdale et al. 2003). From the data we measured $160\mu\text{m}$ fluxes within $64''$ apertures, again using the $24\mu\text{m}$ centroid. The SWIRE $160\mu\text{m}$ data is much shallower than the FIDEL data; however, ECDFS-12514 is detected at the 1.5σ level (a similar detection was made by Dye et al. 2009 using the SWIRE data). ECDFS-4511 is not detected; however, we calculate a 2σ upper limit based on empty aperture fluxes from the SWIRE photometry.

In our SED fitting we use our measured $24\mu\text{m}$, $70\mu\text{m}$ and $160\mu\text{m}$ fluxes, the $250\mu\text{m}$, $350\mu\text{m}$, and $500\mu\text{m}$ fluxes measured by Dye et al. (2009), and the observed $870\mu\text{m}$ fluxes from Weiss et al. (2009). For ECDFS-4511 we do not use the $160\mu\text{m}$ upper limit in the fitting. These fluxes are listed in Table 1.

A joint analysis of the FIDEL $24\mu\text{m}$ and BLAST FIR data by Chary & Pope (2010) has shown that source blending is a problem for many of the BLAST detections. We have used the $24\mu\text{m}$ data to test for contamination of the longer wavelength data of these galaxies by blended sources.

The galaxy ECDFS-12514 is well-isolated on the $24\mu\text{m}$ image. The nearest $24\mu\text{m}$ source is $10''$ away; however, it is extremely faint, only 4% of the flux of the main galaxy. The only nearby bright $24\mu\text{m}$ source is $21''$ away which means if it was luminous at any of the longer wavelengths, excluding $500\mu\text{m}$, it should be resolved as a separate source. That galaxy is only 15% of the main galaxy flux and shows no evidence of a detection in any of the longer wavelength channels, suggesting that the $70 \rightarrow 870\mu\text{m}$ measurements for ECDFS-12514 are probably not contaminated by blended sources.

Galaxy ECDFS-4511 is not completely isolated. There are two well-detected $24\mu\text{m}$ sources within $30''$. The closest is $7''$ away and is 15% of the flux of the main galaxy. The next closest is $11''$ away and is 37% of the flux of the main galaxy. If these companions are luminous in the FIR, neither would be resolved in the FIR data and could cause the FIR flux to be overestimated. Given that at $z \sim 2$, the SED of the main galaxy has been redshifted to where the dust SED peaks in the BLAST bands and the observed $24\mu\text{m}$ /FIR flux ratio is maximal, we expect that the worst-case scenario would be if the nearby galaxies were at the same redshift and had similar FIR SEDs as the main galaxy and therefore contribute equally to the FIR flux. If so, it suggests the FIR fluxes and L_{IR} could be overestimated by as much as $\sim 30\%$. If the companions are at either higher or lower redshift, they are likely to have higher $24\mu\text{m}$ /FIR flux ratios and therefore the overestimate is likely to be less. As we discuss below, this possible overestimate of the L_{IR} of ECDFS-4511 does not change any of the conclusions of this paper.

3. TOTAL INFRARED LUMINOSITIES AND STAR FORMATION RATES

3.1. Spectral Energy Distributions and Fitting

In the top panels of Figure 1 we plot the observed $U \rightarrow 870\mu\text{m}$ flux densities of the galaxies. The $24\mu\text{m} \rightarrow 870\mu\text{m}$ flux densities, which are from dust emission are plotted as black points and the $U \rightarrow 8\mu\text{m}$ flux densities, which are from stellar photospheres are plotted as blue

⁶ <http://irsa.ipac.caltech.edu/data/SPITZER/FIDEL/>

⁷ <http://ssc.spitzer.caltech.edu/mips/dh/>




FIG. 1.— Top panels: U \rightarrow 870 μm SEDs for galaxies ECDFS-4511 (left) and ECDFS-12514 (right). The U \rightarrow 8 μm photometry is plotted in blue, and the best fitting BC03 model to those data is plotted in black. The FIR photometry from Spitzer, BLAST and APEX are plotted in black. The best fitting CE01 model to those data is shown in red. The χ^2 grid for the FIR SEDs is shown in the bottom left panels. The dot dashed line in the upper panel shows the χ^2 of the local template with the same L_{IR} as the best-fit template. Both galaxies are inconsistent with the local template and are best described as "scaled up" versions of lower luminosity galaxies. The dotted FIR SED in the main panel shows the CE01 template that would be chosen based on only the observed 24 μm flux. For these galaxies those templates overestimate the L_{IR} by factors of 3.0 and 6.2, respectively. The bottom right panels show 3" \times 3" NICMOS images of the galaxies.

TABLE 1
MIR/FIR PHOTOMETRIC DATA

Wavelength	ECDFS-4511 flux (mJy)	ECDFS-12514 flux (mJy)	Survey/reference
24 μm	0.57 ± 0.02	1.42 ± 0.03	FIDEL, this paper
70 μm	1.6 ± 1.2	4.2 ± 1.8	FIDEL, this paper
160 μm	< 40	34 ± 20	SWIRE, this paper
250 μm	48 ± 11	35 ± 11	BLAST, Dye et al. (2009)
350 μm	44 ± 9	44 ± 8	BLAST, Dye et al. (2009)
500 μm	19 ± 6	24 ± 6	BLAST, Dye et al. (2009)
870 μm	5.9 ± 1.3	4.7 ± 1.2	LESS, Weiss et al. (2009)

NOTE. — 1) All fluxes are measured fluxes and do not include statistical deboosting corrections. 2) ECDFS-4511 has two 24 μm sources within 11" of the main galaxy. These galaxies cannot be resolved in the FIR data. We estimate that if they have substantial FIR flux they could inflate the FIR fluxes, and L_{IR} by as much as 30% (see text).

points. For reference, the best fit Bruzual & Charlot (2003, hereafter BC03) model to the U \rightarrow 8 μm data from Muzzin et al. (2009) has also been plotted as the solid black line. Figure 1 shows that there is an unambiguous peak in the FIR SEDs of the galaxies at $\sim 300\mu\text{m}$, a clear signature of the redshifted thermal emission from dust.

We fit the 24 μm \rightarrow 870 μm data using the models of Chary & Elbaz (2001; hereafter CE01). The CE01 models are a grid of 105 synthetic SEDs that were constructed to reproduce the correlations between the observed MIR

and FIR fluxes of local galaxies. Each model SED is associated with an L_{IR} ranging between $3 \times 10^8 L_\odot$ and $6 \times 10^{13} L_\odot$. We fit the 24 μm \rightarrow 870 μm data using each of the 105 template SEDs using a χ^2 minimization method where the amplitude of the SED is the only free parameter.

The galaxy ECDFS-4511 is the most FIR-luminous of the two galaxies and is well-fit by the SED template of a ULIRG. In the CE01 models that template is associated with an $L_{\text{IR}} = 1.1 \times 10^{12} L_\odot$; however, ECDFS-4511 has

$L_{IR} = 8.9^{+1.1}_{-1.4} \times 10^{12} L_{\odot}$, eight times larger than the luminosity of the local template.

The second galaxy, ECDFS-12514, is three times brighter at $24\mu\text{m}$ than ECDFS-4511; however, it has a lower L_{IR} . The best-fit template is a LIRG template with strong PAH features and is associated with $L_{IR} = 1.2 \times 10^{11} L_{\odot}$. The L_{IR} of ECDFS-12514 is $5.7^{+1.2}_{-0.1} \times 10^{12} L_{\odot}$, almost 50 times more luminous than the local template. If the local template is correct, the large $24\mu\text{m}$ flux from ECDFS-12514 likely arises from strong rest-frame $7.8\mu\text{m}$ PAH emission redshifted into the observed $24\mu\text{m}$ bandpass.

The key result in Figure 1 is that although the $z = 2$ galaxies have FIR SEDs that are similar to local FIR luminous galaxies, their luminosities are roughly an order of magnitude larger. It appears that both galaxies are strongly "scaled up" versions of much lower luminosity galaxies in the local universe. The dot-dashed line on the χ^2 grids in Figure 1 show the χ^2 of the fit to the local template that is associated with the best fit L_{IR} of ECDFS-4511 and ECDFS-12514. Those χ^2 s are significantly worse than the best-fit χ^2 and show that the SEDs of the $z \sim 2$ galaxies are inconsistent with having the same SED shape of local templates of similar luminosity at $> 25\sigma$.

3.2. Comparison of FIR and $24\mu\text{m}$ Star Formation Rates

Direct constraints on the cool dust SED allow us to address the issue of how well L_{IR} , and $\text{SFR}(L_{IR})$ can be determined for $z \sim 2$ galaxies using $24\mu\text{m}$ data alone. The most common practice for determining L_{IR} from $24\mu\text{m}$ photometry is to select the local template with an L_{IR} that would match the observed $24\mu\text{m}$ flux if redshifted to the redshift of the source galaxy. In Figure 1 we plot the CE01 template that would be selected based on using that method as the red dotted line.

The extraordinary $24\mu\text{m}$ flux of both galaxies requires that they use the most luminous templates in the CE01 library. The implied L_{IR} for ECDFS-4511 and ECDFS-12514 is $2.8 \times 10^{13} L_{\odot}$ and $3.8 \times 10^{13} L_{\odot}$, respectively. These templates overestimate the L_{IR} from the full SED fit by factors of 3.0 and 6.2 respectively, and it is clear from Figure 1 that they cannot simultaneously fit both the $24\mu\text{m}$ flux densities and the FIR flux densities. If these galaxies are representative of most FIR-luminous galaxies in the distant universe, it suggests we may be systematically overestimating total SFRs by similar factors.

Similar results have also been found by Papovich et al. (2007) and Murphy et al. (2009). Papovich et al. (2007) compared L_{IR} determined from $24\mu\text{m}$ data to that determined using the combined $24\mu\text{m}$, $70\mu\text{m}$, and $160\mu\text{m}$ photometry and found that for the most luminous galaxies ($S_{24} > 250\mu\text{Jy}$) the $24\mu\text{m}$ photometry overestimated the L_{IR} by factors of 2 – 10. Using a sample of 14 MIR luminous galaxies at $1.4 < z < 2.6$ with Spitzer IRS spectra and SCUBA $850\mu\text{m}$ photometry Murphy et al. (2009) found that $24\mu\text{m}$ fluxes combined with local templates systematically overestimated the L_{IR} by a factor of ~ 5 .

3.3. Comparison of FIR and $\text{H}\alpha$ Star Formation Rates




FIG. 2.— A_v determined from the Balmer decrement vs. A_v determined from a fit to the galaxy stellar population for the 12 emission line galaxies from Kriek et al. (2007), Kriek et al. (2009a), and van Dokkum et al. (2005) with both $\text{H}\beta$ and $\text{H}\alpha$ within the NIR window. Only two galaxies have secure $\text{H}\beta$ detections, the remainder are plotted as 1σ lower limits. The two BLAST-detected galaxies are indicated as the red lower limits. The dotted line shows the one-to-one relation, and the solid line is the relation $E(B-V)_{\text{stellar}} = 0.44E(B-V)_{\text{nebular}}$ suggested by Calzetti et al. (1997). The two galaxies with $\text{H}\beta$ detections are in reasonable agreement with the Calzetti relation. All of the galaxies with lower limits are consistent with the Calzetti relation but only about half are consistent with the one-to-one relation.

The FIR-derived SFRs can also be compared to the dust corrected $\text{H}\alpha$ SFRs for these galaxies. $\text{H}\alpha$ equivalent widths (EWs) were measured for both galaxies using moderate-resolution NIR spectra from SINFONI by Kriek et al. (2007). We convert their $\text{EW}(\text{H}\alpha)$ to an $\text{H}\alpha$ line flux by multiplying by the continuum flux at 6563\AA estimated using the BC03 SED model fits from Muzzin et al. (2009).

We assess the extinction for the $\text{H}\alpha$ flux using both the fit to the stellar SED ($U \rightarrow 8\mu\text{m}$) and the Balmer decrement. For the SED fits we use the Muzzin et al. (2009) best-fit value of A_v , which is derived from fitting BC03 models using solar metallicity and the Calzetti et al. (2000) dust law (hereafter $A_{v,SED}$). For the A_v from the Balmer decrement (hereafter $A_{v,Balmer}$) we assume the Calzetti et al. (2000) dust law and an intrinsic ratio of $\text{H}\alpha/\text{H}\beta = 2.86$. Although both galaxies have high S/N $\text{H}\alpha$ detections in the SINFONI spectra, neither has a secure $\text{H}\beta$ detection. Therefore instead of a true Balmer decrement we calculate 1σ lower limits⁸ using the 1σ upper limits on $\text{H}\beta$ determined by Kriek et al. (2007). The $\text{H}\beta$ upper limits have been corrected for absorption by the stellar population using a BC03 SED fit to the broad-band photometry and the NIR spectroscopy.

In Figure 2 we plot $A_{v,Balmer}$ vs. the $A_{v,SED}$ for the two galaxies as the red lower limits. For comparison, measurements of the 9 other $z \sim 2$ galaxies in Kriek et al. (2007), and the galaxy from Kriek et al. (2009b) are also plotted. Two of these galaxies (ID#’s 1030-807 and CDFS-6202) have secure $\text{H}\beta$ detections, the remainder are plotted as 1σ lower limits. The one-to-one relation is shown as the dotted line.

⁸ Upper limits on $\text{H}\beta$ translate to lower limits on $\text{H}\alpha/\text{H}\beta$ and lower limits on $A_{v,Balmer}$

The two galaxies with $H\beta$ detections do not agree well with the one-to-one relation. The lower limits for approximately half the galaxies are also inconsistent with the one-to-one relation. A similar disagreement amongst A_v measurements was found for local galaxies by Calzetti (1997). She showed that this difference can be understood as greater extinction towards the star forming regions, where the $H\alpha$ originates, than to the overall stellar population. The Calzetti (1997) relation for local galaxies is $E(B-V)_{stars} = (0.44 \pm 0.03)E(B-V)_{nebular}$. In Figure 2 we plot that relation converted to A_v assuming $R_v = 4.05$ (Calzetti et al. 2000) as the solid line. The Calzetti (1997) relation seems to provide a better description of the overall population of $z \sim 2$ star forming galaxies, albeit that most galaxies only have lower limits.

Given the uncertainty in the $A_{v,Balmer}$, and the well-determined $A_{v,SED}$ we have chosen to dust correct the $H\alpha$ flux using the $A_{v,SED}$ but using the $E(B-V)_{stars} = (0.44 \pm 0.03)E(B-V)_{nebular}$ relation. We convert this to a SFR using the Kennicutt (1998) relation, $SFR(H\alpha) = 0.57 \cdot L(H\alpha)/1.26 \times 10^{-41} \text{ ergs s}^{-1}$, where the factor of 0.57 has been added to convert from a Salpeter IMF to a Chabrier IMF. Applying these conversions we find that ECDFS-4511 and ECDFS-12514 have $SFR(H\alpha) = 620^{+260}_{-140} M_\odot \text{ yr}^{-1}$ and $1210^{+750}_{-40} M_\odot \text{ yr}^{-1}$, respectively. We note that the majority of the error budget for these measurements comes from the uncertainty in the extinction correction, not from random errors in the measurement of the $H\alpha$ line flux.

We convert the L_{IR} to a SFR using the Kennicutt (1998) relation, $SFR(L_{IR}) = 0.57 \cdot L_{IR}/5.8 \times 10^9 L_\odot$, where again the factor of 0.57 is to convert to a Chabrier IMF. We derive $SFR(L_{IR}) = 870^{+150}_{-140} M_\odot \text{ yr}^{-1}$ and $560^{+120}_{-10} M_\odot \text{ yr}^{-1}$ for ECDFS-4511 and ECDFS-12514, respectively.

Comparing the two SFR indicators shows that they agree to within roughly a factor of 2 for both galaxies. Such an agreement is good considering that there are still systematic uncertainties in the conversion of both L_{IR} and $L(H\alpha)$ to a SFR that are of that order (Kennicutt et al. 2009).

It is worth noting that the agreement in SFRs only occurs because L_{IR} is determined from the full FIR SED fits, and because the extra extinction correction has been applied to the $L(H\alpha)$. If we do not use the L_{IR} from the FIR SED fit and instead use the $SFR(24\mu\text{m})$ we find that the IR SFR is larger than the $SFR(H\alpha)$ by factors of 4.8 and 3.4 for ECDFS-4511 and ECDFS-12514, respectively. Likewise, if we extinction correct the $L(H\alpha)$ using the $A_{v,SED}$ without extra extinction we find that the $SFR(L_{IR})$ is larger than the $SFR(H\alpha)$ by a factor of 5.4 and 2.5; however, this time because the $SFR(H\alpha)$ is underestimated. If we compare the $SFR(24\mu\text{m})$ to the $SFR(H\alpha)_{SED}$, the $SFR(24\mu\text{m})$ is larger by factors of 18.7, and 18.6.

4. ARE THE BLAST-DETECTED GALAXIES TYPICAL OF $z \sim 2$ MASSIVE GALAXIES?

The reasonable agreement between the FIR-derived and $H\alpha$ derived SFRs in the previous section shows that provided sufficient data to robustly determine both L_{IR} and a dust corrected $L(H\alpha)$, the locally-calibrated conversions of $L_{IR} \rightarrow \text{SFR}$ and $L(H\alpha) \rightarrow \text{SFR}$ appear to




FIG. 3.— Top Panel: SFR derived from the $24\mu\text{m}$ fluxes and the luminosity dependent CE01 templates vs. the SFR determined from the dust corrected $L(H\alpha)$ for the full sample of K-selected galaxies in Kriek et al. (2008). The luminosity dependent templates systematically overestimate the $SFR(H\alpha)$ for the majority of galaxies. Bottom Panel: Log-average of the SFR derived from the $24\mu\text{m}$ fluxes and all 105 CE01 templates vs. $SFR(H\alpha)$. An average of the local templates provides IR SFRs that are in better agreement with the $SFR(H\alpha)$. The luminosity dependent method requires that the majority of galaxies use the most luminous templates, whereas the template averaging forces the L_{IR} to come from cooler templates. The better agreement from cooler templates suggests the entire sample of massive $z \sim 2$ galaxies would be consistent with being “scaled up” versions of lower luminosity galaxies.

provide consistent results for these galaxies. Before we use these two galaxies to draw conclusions about the star formation properties of the larger population of $z \sim 2$ massive galaxies, we must consider if they are representative. With $24\mu\text{m}$ flux densities of $570 \pm 18 \mu\text{Jy}$ and $1417 \pm 28 \mu\text{Jy}$, these two galaxies are the brightest $24\mu\text{m}$ sources in the entire Kriek et al. (2008) sample. That they are the only ones detected in both the BLAST and APEX observations of the ECDFS is potentially a strong selection effect.

One way to test how representative the BLAST-detected galaxies are of the entire K-selected sample is to compare the $SFR(24\mu\text{m})$ and $SFR(H\alpha)$ of the full sample and see if it shows the same disagreement as the BLAST-detected galaxies. MIPS $24\mu\text{m}$ observations of the MUSYC fields were obtained as part of Spitzer program GO-30873 (PI: Labbé). We reduced that data with the standard MOPEX procedures and derived $24\mu\text{m}$ fluxes for galaxies in the Kriek et al. (2008) sample using the method described in § 2. There are 24 galaxies in the Kriek et al. sample where $H\alpha$ is observable in the NIR window from the ground, and of these galaxies 15 are detected at $24\mu\text{m}$. The MUSYC $24\mu\text{m}$ data is much shallower than the FIDEL data, so the non-detection of some galaxies, depending on their SFR and redshift is

expected. For the undetected galaxies we calculate 5σ upper limits based on empty aperture fluxes in the $24\mu\text{m}$ mosaics.

In the top panel of Figure 3, we plot the $\text{SFR}(24\mu\text{m})$ vs. the $\text{SFR}(\text{H}\alpha)$ for these galaxies. Galaxies that are candidates for hosting an AGN from either an X-ray detection, emission-line ratios (Kriek et al. 2007) or for having a powerlaw SED in the IRAC bands (Muzzin et al. 2009) are plotted as red symbols (this includes both BLAST-detected galaxies, see § 5.3). For reference, the BLAST-detected galaxies are plotted as stars at their SED-fit $\text{SFR}(\text{LIR})$ locations with arrows connecting them to their $\text{SFR}(24\mu\text{m})$. Just like the BLAST sources, the majority of galaxies have a $\text{SFR}(24\mu\text{m})$ that is systematically larger than the $\text{SFR}(\text{H}\alpha)$. This suggests that the local luminosity dependent templates probably overestimate the IR SFR of the majority of $z \sim 2$ massive galaxies.

Franx et al. (2008) and Wuyts et al. (2009) also noticed that the SFRs of $z \sim 2$ galaxies tended to be larger when inferred from $24\mu\text{m}$ flux and the local luminosity-dependent templates than the SFR determined using a fit to the stellar SED ($\text{SFR}(\text{SED})$). They advocated using the LIR taken from the log-average of the local templates and argued that this produces SFRs that are in better-agreement with the $\text{SFR}(\text{SED})$. For comparison, we determine the $\text{SFR}(24\mu\text{m})$ using that method (although we use the CE01 templates, whereas they use the Dale & Helou 2002 templates), and compare with the $\text{SFR}(\text{H}\alpha)$ in the bottom panel of Fig 3.

Clearly the agreement between SFRs is much better when $\text{SFR}(24\mu\text{m})$ is determined using template averaging method. Given that when using the luminosity dependent templates, most of our galaxies require the most luminous templates to match the $24\mu\text{m}$ flux, it suggests the better agreement between SFRs with the template averaging method is because it includes numerous lower luminosity templates in the average. This suggests that the larger sample of K-selected galaxies, like the BLAST-detected galaxies, can be described as scaled up versions of lower luminosity galaxies. Furthermore, it appears that despite their exceptionally large FIR luminosities, the cooler SEDs of the BLAST-detected galaxies may be representative of the larger population of massive galaxies at $z \sim 2$.

5. DISCUSSION

5.1. Dust Temperatures and Evidence for "Scaled Up" Cool Galaxies

It is remarkable that despite an $\text{LIR} \sim 10^{13} L_{\odot}$, both of the BLAST-detected $z \sim 2$ galaxies are better fit with local templates that have PAH features and a cool "cold" dust bump. In the nearby universe, galaxies with $\text{LIR} \sim 10^{13} L_{\odot}$ tend to have both the warmest "cold" dust ($T_{\text{dust}} \sim 10 - 70\text{K}$, emission peaking in the FIR) and significant amounts of "hot" dust ($T_{\text{dust}} \sim 70 - 500\text{K}$, emission peaking in the MIR). Although both of these dust components can come from deeply embedded star formation (e.g., Tran et al. 2001), it is thought that the dominant radiation source in most, if not all nearby galaxies with $L \sim 10^{13} L_{\odot}$ is an AGN (e.g., Genzel et al. 1998; Lutz et al. 1998).

By contrast, the best fit templates of the $z \sim 2$ galaxies

are much cooler ULIRG and LIRG templates which are most frequently associated with star forming galaxies in the local universe (e.g., Armus 2009). These templates also tend to have stronger PAH features that are more prevalent because the "hot" dust component from the AGN no longer dominates the MIR emission.

The first claims that IR luminous galaxies at $z \sim 2$ may be scaled up versions of lower luminosity cooler galaxies came from $24\mu\text{m}$ observations of submillimeter-selected galaxies (SMGs). Pope et al. (2006) showed that the SEDs of SMGs peak at longer wavelengths than expected based on an extrapolation of their $24\mu\text{m}$ flux using local templates, arguing that SMGs are cooler at a given luminosity than the local templates. Subsequent MIR spectroscopic observations of SMGs from *Spitzer* by Lutz et al. (2005), Pope et al. (2008), and Menendez-Delmestre et al. (2009) all found strong PAH features and relatively weak MIR continuum in the SMGs suggesting much less "hot" dust than is usually found in local galaxies of comparable LIR . Similarly, Papovich et al. (2009) found a weaker than expected rest-frame $24\mu\text{m}$ luminosity for a lensed SMG, arguing for the lack of a "hot" dust component, and hence a scaled up cool galaxy.

With access to the FIR we can determine a dust temperature (T_d) independent of the MIR SED and compare this to the local T_d - LIR relation. We fit the $160\mu\text{m} \rightarrow 870\mu\text{m}$ data ($50\mu\text{m} \rightarrow 300\mu\text{m}$ rest-frame) to modified blackbody curves of the form $S(\nu) = A\nu^{\beta}B(\nu, T_d)$, where A is a normalization, B is the Planck function, and β accounts for frequency-dependent dust emissivity. Even with detections in numerous bands the S/N of the data is too low to constrain both β and T_d simultaneously so we have assumed $\beta = 1.5$, a typical value for submillimeter galaxies. The $70\mu\text{m}$ points are omitted in the fitting because they correspond to rest-frame $23\mu\text{m}$ and are too far into the MIR to include in a single-component blackbody fit. We note that including the $70\mu\text{m}$ data actually produces identical best-fit temperatures; however, the χ^2 of fits are substantially larger because no single-temperature model can describe the $23\mu\text{m} \rightarrow 300\mu\text{m}$ rest-frame simultaneously.

We find dust temperatures of 40^{+2}_{-1}K and 41^{+5}_{-7}K for ECDFS-4511 and ECDFS-12514, respectively. These correspond to rest-frame monochromatic IRAS color measurements, $C \equiv \log_{10}(f_{60\mu\text{m}}/f_{100\mu\text{m}})$, of $-0.05^{+0.05}_{-0.03}$ and $-0.03^{+0.09}_{-0.20}$, respectively. Comparing these to the C- LIR relation measured by Chapin et al. (2009) in the nearby universe using IRAS shows that these galaxies have temperatures typical of nearby galaxies with $\text{LIR} = 10^{11} - 10^{12} L_{\odot}$. They are also similar to the T_d of local LIRGS/ULIRGS measured by Clements et al. (2010). Therefore, with the combined *Spitzer*, BLAST and APEX data we can now confirm that some massive galaxies at $z \sim 2$ not only have PAH-dominated MIR SEDs similar to lower luminosity local galaxies, but they also have "cold" dust SEDs consistent with being scaled up versions of lower luminosity galaxies. This suggests that star formation may be occurring in a different environment within galaxies at $z \sim 2$ compared to $z \sim 0$.

Local galaxies with SFRs $> 100 M_{\odot} \text{ yr}^{-1}$ tend to have compact nuclear starbursts which are the cause of the warmer dust temperatures (e.g., Sanders & Mirabel 1996). The $z \sim 2$ galaxies have SFRs $\sim 1000 M_{\odot} \text{ yr}^{-1}$,

but similar dust temperatures which suggests that the star formation may be occurring in multiple large starburst regions throughout the galaxy, rather than concentrated in a single compact starburst which would presumably have a substantially hotter dust temperature. Indeed, a lensed $z > 2$ galaxy with precisely such characteristics has recently been observed by Swinbank et al. (2010a) using interferometric submillimeter observations.

Kriek et al. (2009a) presented SINFONI-IFU $H\alpha$ spectra of both these galaxies. Their maps show that the $H\alpha$ emission is resolved on scales of a few kpc, and is also consistent with the extended star formation hypothesis. The greater availability of gas at high redshift may permit more spatially extended star formation at a high rate, something that is uncommon in local galaxies.

5.2. Morphologies

In the inset of Figure 1 we show $3'' \times 3''$ NIC2 F160W images of the galaxies from Kriek et al. (2009a). At $z \sim 2$ these correspond to rest-frame V-band morphologies and span an angular size of ~ 25 kpc. Like most of the high redshift ULIRGs and HLIRGs previously observed with HST (e.g., Chapman et al. 2003; Dasyra et al. 2008; Swinbank et al. 2010b) the galaxies clearly have disturbed morphologies. Kriek et al. (2009a) measure effective radii (R_e) of 5 and 3 kpc for ECDFS-4511 and ECDFS-12514, respectively, which means that both galaxies are comparable in size to local galaxies of similar mass. Rest-frame optical morphologies make it difficult to identify the location of the star formation (which is presumably highly obscured); however, the disturbed nature of both galaxies shows that these are complex systems, possibly "stream-fed" galaxies (Dekel et al. 2009) or mergers. Unlike the FIR SEDs, the morphologies do not appear to be scaled up versions of local star forming disk galaxies.

5.3. AGN Contribution to the MIR and FIR Emission

Despite SEDs that are associated with local star forming galaxies, part of the MIR and FIR emission in both galaxies almost certainly comes from an AGN. Using emission line ratios and the BPT diagram (Baldwin et al. 1981), Kriek et al. (2007) found that both galaxies were in the region of the diagram that Kewley et al. (2006) considered to have "composite" spectra with contributions from both star formation and an AGN.

ECDFS-4511 is detected in the 2 Ms *Chandra* map of the CDFS (Luo et al. 2008), and has an $L_x \sim 2 \times 10^{44}$ erg s $^{-1}$, implying it hosts a Seyfert-luminosity AGN. ECDFS-12514 is located in the 250 ks map of the ECDFS (Virani et al. 2006), but is undetected. Although undetected in the X-rays, ECDFS-12514 is formally classified as an IRAC powerlaw galaxy ($F_\nu \propto \nu^\alpha$, $\alpha < -0.5$; e.g., Donley et al. 2007, see Figure 1) which makes it a candidate for an obscured AGN.

Without more data we cannot constrain the precise fraction of the MIR and FIR light that comes from the AGN; however, the consistency of the MIR/FIR SED with local templates that are star formation dominated, and the "composite" emission line ratios suggest that young stars are probably the dominant energy source heating the dust. Pope et al. (2008) and Menendez-Delmestre et al. (2009) showed that the MIR SEDs of

most SMGs (the BLAST-detected galaxies are formally SMGs) are still dominated by PAHs and that AGN only contribute $\sim 30\%$ of the total MIR emission. Putting the evidence together the most plausible scenario is that both galaxies host a moderate-luminosity AGN, but the AGN does not dominate the MIR/FIR SED. Nevertheless, we note that both our $SFR(L_{IR})$ and $SFR(H\alpha)$ should be formally considered upper limits as the AGN will contribute to both of these.

5.4. Implications for SFRs at $z \sim 2$ and MIR-excess Galaxies

The results of our FIR SED fitting show that caution is required when determining and interpreting the SFRs of $z \sim 2$ galaxies based on limited data. Using only $24\mu\text{m}$ photometry to infer SFRs at $z \sim 2$ may work well for lower luminosity, or less obscured galaxies (e.g., Papovich et al. 2007; Reddy et al. 2010); however, the local templates can overestimate the $SFR(L_{IR})$ of the most FIR luminous galaxies by as much as a factor of ~ 6 (see also Papovich et al. 2007; Murphy et al. 2009). Likewise, the Balmer decrement measurements for these galaxies seem to indicate extra extinction toward the star forming regions. It appears that on average this relation is similar to that suggested by Calzetti et al. (1997); however, the lack of secure $H\beta$ detections in our sample precludes quantitative statements. To correctly determine the $SFR(H\alpha)$ of individual galaxies, rather than ensemble averages requires high quality Balmer decrement measurements.

The extra extinction towards the star forming regions suggested by the Balmer decrements also indicates that the UV SFRs for some of these IR luminous galaxies may not be reliable because the A_v is large enough that parts of the star forming regions are optically-thick. The SFRs estimated from a fit to the stellar populations ($SFR(SED)$; effectively a dust corrected UV SFR) by Muzzin et al. (2009) for ECDFS-4511 and ECDFS-12514 are $168^{+41}_{-63} M_\odot \text{ yr}^{-1}$ and $145^{+160}_{-0} M_\odot \text{ yr}^{-1}$, and underestimate the $SFR(L_{IR})$ of the galaxies by factors of 5.2 and 3.5, respectively. If we compare the $SFR(SED)$ to the $SFR(24\mu\text{m})$ as in e.g., Daddi et al. (2007b) we find that these galaxies have MIR SFRs 18 – 25 times greater than the UV SFRs, and are extremely strong "MIR excess" galaxies.

The purpose of the Daddi et al. (2007b) classification was to identify candidate obscured AGN, and given that it is likely that both galaxies contain an AGN it appears that method successfully identifies such sources. However, MIR excesses of factors of 18 – 25 would suggest that the AGN is the dominant MIR energy source for both galaxies, something that is unlikely given the shape of the MIR/FIR SEDs and the emission line ratios. This calls into question whether galaxies with weaker inferred MIR excesses, such those near the cutoff of $SFR(UV_{corr})/SFR(MIR+UV) \sim 3$ suggested by Daddi et al. (2007b) have a significant contribution from an AGN.

5.5. Comparison to Models

Our results suggest that a substantial population of massive $z \sim 2$ galaxies are "scaled up" versions of lower luminosity galaxies. Clearly such a population needs to

be understood within the context of modern galaxy formation and evolution models. Recently, Hopkins et al. (2010) have used their merger-driven evolutionary model to predict the IR luminosity functions of galaxies and quasars between $0 < z < 6$. In their model they argue that there is a threshold in L_{IR} above which disks, merger-driven star formation, and obscured AGN are the dominant IR energy source of galaxies. At $z = 0$ these limits are $\log(L_{IR}) \geq 11.5$ for merger-driven star formation, and $\log(L_{IR}) \geq 12.5$ for obscured AGN. Based on the evolution of the merger rate and gas fraction to $z \sim 2$ they suggest that these thresholds evolve in $\log(L_{IR})$ to $\log(L_{IR}) \geq 12.5$ for merger-driven star formation and $\log(L_{IR}) \geq 13.0$ for obscured AGN. With $\log(L_{IR}) \sim 13.0$, the BLAST galaxies lie at an L_{IR} that is near the suggested transition point between merger-driven star formation dominated or AGN dominated.

Interestingly, the FIR SEDs of these galaxies do not look like those of the AGN-dominated HLIRGs detected by *ISO* (Genzel et al. 1998), or of the average nearby ULIRG, which are mostly merger-driven starbursts (e.g., Sanders et al. 1998). Instead, these galaxies look like scaled up versions of even lower luminosity systems, which are not universally associated with mergers or AGN. Taken at face value, it appears they do not fit well within the Hopkins et al. (2010) description.

On the other hand, the NIC2 morphologies of the galaxies are clearly disturbed enough to be considered candidates for undergoing mergers (Kriek et al. 2009b). If these galaxies are merger-driven starbursts, it may be that they are missing the centrally-concentrated starburst which produces the much hotter dust seen in these galaxies in the local universe. As Hopkins et al. (2010) point out, the efficiency of a merger at funneling gas and stars into the core of the galaxy is proportional to the gas fraction, with higher gas fraction mergers being less efficient. If both galaxies have high gas fractions most of the star formation could be occurring on much larger physical scales than similar mass starbursts in the local universe, hence the cool dust SED.

Galaxies with similar stellar masses ($M \sim 10^{11} M_{\odot}$) and gas fractions as high as 80% have been observed in the distant universe (e.g., Tacconi et al. 2010) and could be analogues of the BLAST-detected galaxies. If that is the case, it begs the question of what are the descendants of extended, massive, gas-rich star forming galaxies at $z \sim 2$? Local spheroids seem like an obvious candidate; however, some fraction of the progenitors of those systems are clearly small and quiescent at $z \sim 2$ (e.g., van Dokkum et al. 2010; Hopkins et al. 2010; Bezanson et al. 2009; van der Wel et al. 2008). If the large and active BLAST-sources are also progenitors of massive local spheroids it suggests there may be very different evolutionary paths which arrive at the same galaxies.

These galaxies also appear to be consistent with the "stream-fed" galaxies proposed by Dekel et al. (2009) and the scaled up "turbulent disks" discussed by Genzel et al. (2008). In their model, Dekel et al. (2009) suggest that massive galaxies may grow quickly through rapid cold gas accretion and clumpy star formation which funnels stars into a central bulge. Given that these cold flows often separate into multiple large star forming clumps, it might be expected that these clumps produce cooler dust

temperatures at fixed SFR than centrally concentrated starbursts.

The SFRs of the BLAST galaxies are higher than the $100 - 200 M_{\odot} \text{ yr}^{-1}$ predicted for "stream-fed" galaxies by Dekel et al. (2009). The BLAST galaxies are the most IR luminous of the K-selected sample of Kriek et al. (2008), so if they are "stream-fed" galaxies, they may just represent the most active tail of the distribution.

High quality kinematic data could potentially distinguish between merger-dominated or accretion-dominated systems. These galaxies do have SINFONI IFU observations; however, these data lack the S/N to reliably determine V/σ . Therefore, at present based on the MIR/FIR properties and morphologies we cannot distinguish between evolutionary models such suggested by both Hopkins et al. (2010) and Dekel et al. (2009).

6. SUMMARY

Using a combination of data from the Spitzer, BLAST and APEX observatories we have presented some of the first well-sampled FIR SEDs for two $z \sim 2$ galaxies with rest-frame optical spectroscopy. In both galaxies we see a clear peak in the FIR emission at $\sim 300 \mu\text{m}$ which is caused by the redshifted thermal emission from dust. The FIR SEDs of the galaxies are well described by local templates; however, the L_{IR} appears to be an order of magnitude larger than the L_{IR} associated with the locally calibrated templates. This suggests that these galaxies are likely to be scaled up versions of "cool" local galaxies, rather than analogues of nearby HLIRG templates.

A comparison of $\text{SFR}(L_{IR})$ and $\text{SFR}(H\alpha)$ for these two galaxies shows that they are consistent; however, if we use only the $24 \mu\text{m}$ photometry and the local luminosity dependent templates to determine the SFR we find that this overestimates the $\text{SFR}(H\alpha)$ by as much as a factor of 6. A larger sample of 24 galaxies from the Kriek et al. GNIRS sample shows the same relation between $\text{SFR}(24 \mu\text{m})$ and $\text{SFR}(H\alpha)$ and suggests that many massive galaxies at $z \sim 2$ are likely to be scaled up versions of lower luminosity galaxies.

Although based on just a handful of galaxies, our results show that FIR luminous galaxies at $z \sim 2$ are clearly different than their counterparts in the local universe. In the coming years the *Herschel* and JWST observatories will provide us with large amounts of high-quality FIR observations on the population of luminous galaxies at $z > \sim 2$. Likewise, upcoming NIR multi-object spectrograph will provide better $H\alpha$ and $H\beta$ measurements for those galaxies, and the ALMA observatory will provide spatially-resolved FIR maps of these galaxies. This plethora of new, high-quality data will allow us to better understand how the population of luminous, "cool" star forming galaxies fit into the overall picture of galaxy evolution.

We thank the anonymous referee whose comments helped improve the quality of the paper. Support for program HST-GO-11135.08 was provided by NASA through a grant from the Space Telescope Science Institute.

REFERENCES

- Armus, L. 2009, *Astronomical Society of the Pacific Conference Series*, 408, 105
- Baldwin, J. A., Phillips, M. M., & Terlevich, R. 1981, *PASP*, 93, 5
- Bertin, E., & Arnouts, S. 1996, *A&AS*, 117, 393
- Bezanson, R., van Dokkum, P. G., Tal, T., Marchesini, D., Kriek, M., Franx, M., & Coppi, P. 2009, *ApJ*, 697, 1290
- Bruzual, G., & Charlot, S. 2003, *MNRAS*, 344, 1000
- Buat, V., et al. 2007, *ApJS*, 173, 404
- Buat, V., et al. 2005, *ApJ*, 619, L51
- Calzetti, D., Armus, L., Bohlin, R. C., Kinney, A. L., Koornneef, J., & Storch-Bergmann, T. 2000, *ApJ*, 533, 682
- Calzetti, D. 1997, *AJ*, 113, 162
- Caputi, K. I., et al. 2007, *ApJ*, 660, 97
- Chapman, S. C., Windhorst, R., Odewahn, S., Yan, H., & Conselice, C. 2003, *ApJ*, 599, 92
- Chary, R.-R., & Pope, A. 2010, *arXiv:1003.1731*
- Chary, R., & Elbaz, D. 2001, *ApJ*, 556, 562
- Clements, D. L., Dunne, L., & Eales, S. 2010, *MNRAS*, 403, 274
- Daddi, E., et al. 2007, *ApJ*, 670, 173
- Daddi, E., et al. 2007, *ApJ*, 670, 156
- Dale, D. A., & Helou, G. 2002, *ApJ*, 576, 159
- Damen, M., Labbé, I., Franx, M., van Dokkum, P. G., Taylor, E. N., & Gawiser, E. J. 2009, *ApJ*, 690, 937
- Dasyra, K. M., Yan, L., Helou, G., Surace, J., Sajina, A., & Colbert, J. 2008, *ApJ*, 680, 232
- Dekel, A., et al. 2009, *Nature*, 457, 451
- Devlin, M. J., et al. 2009, *Nature*, 458, 737
- Donley, J. L., Rieke, G. H., Pérez-González, P. G., Rigby, J. R., & Alonso-Herrero, A. 2007, *ApJ*, 660, 167
- Dye, S., et al. 2009, *ApJ*, 703, 285
- Elbaz, D., Cesarsky, C. J., Chanial, P., Aussel, H., Franceschini, A., Fadda, D., & Chary, R. R. 2002, *A&A*, 384, 848
- Franx, M., van Dokkum, P. G., Schreiber, N. M. F., Wuyts, S., Labbé, I., & Toft, S. 2008, *ApJ*, 688, 770
- Genzel, R., et al. 2008, *ApJ*, 687, 59
- Genzel, R., et al. 1998, *ApJ*, 498, 579
- Hopkins, A. M., Connolly, A. J., Haarsma, D. B., & Cram, L. E. 2001, *AJ*, 122, 288
- Hopkins, P. F., Younger, J. D., Hayward, C. C., Narayanan, D., & Hernquist, L. 2010, *MNRAS*, 402, 1693
- Juneau, S., et al. 2005, *ApJ*, 619, L135
- Kartaltepe, J. S., et al. 2010, *ApJ*, 709, 572
- Kennicutt, R. C., et al. 2009, *ApJ*, 703, 1672
- Kennicutt, R. C., Jr. 1998, *ApJ*, 498, 541
- Kriek, M., van Dokkum, P. G., Franx, M., Illingworth, G. D., & Magee, D. K. 2009, *ApJ*, 705, L71
- Kriek, M., van Dokkum, P. G., Labbé, I., Franx, M., Illingworth, G. D., Marchesini, D., & Quadri, R. F. 2009, *ApJ*, 700, 221
- Kriek, M., et al. 2008, *ApJ*, 677, 219
- Kriek, M., et al. 2007, *ApJ*, 669, 776
- Le Floc'h, E., et al. 2005, *ApJ*, 632, 169
- Lonsdale, C. J., et al. 2003, *PASP*, 115, 897
- Luo, B., et al. 2008, *ApJS*, 179, 19
- Lutz, D., Valiante, E., Sturm, E., Genzel, R., Tacconi, L. J., Lehnert, M. D., Sternberg, A., & Baker, A. J. 2005, *ApJ*, 625, L83
- Lutz, D., Spoon, H. W. W., Rigopoulou, D., Moorwood, A. F. M., & Genzel, R. 1998, *ApJ*, 505, L103
- Magnelli, B., Elbaz, D., Chary, R. R., Dickinson, M., Le Borgne, D., Frayer, D. T., & Willmer, C. N. A. 2009, *A&A*, 496, 57
- Marillac, D., Elbaz, D., Chary, R. R., Dickinson, M., Galliano, F., & Morrison, G. 2006, *A&A*, 451, 57
- Menéndez-Delmestre, K., et al. 2009, *ApJ*, 699, 667
- Murphy, E. J., Chary, R.-R., Alexander, D. M., Dickinson, M., Magnelli, B., Morrison, G., Pope, A., & Teplitz, H. I. 2009, *ApJ*, 698, 1380
- Muzzin, A., Marchesini, D., van Dokkum, P. G., Labbé, I., Kriek, M., & Franx, M. 2009, *ApJ*, 701, 1839
- Papovich, C., Rudnick, G., Rigby, J. R., Willmer, C. N. A., Smith, J.-D. T., Finkelstein, S. L., Egami, E., & Rieke, M. 2009, *ApJ*, 704, 1506
- Papovich, C., et al. 2007, *ApJ*, 668, 45
- Pérez-González, P. G., et al. 2005, *ApJ*, 630, 82
- Pope, A., et al. 2008, *ApJ*, 675, 1171
- Pope, A., et al. 2006, *MNRAS*, 370, 1185
- Reddy, N. A., Erb, D. K., Pettini, M., Steidel, C. C., & Shapley, A. E. 2010, *arXiv:1002.0837*
- Reddy, N. A., Steidel, C. C., Fadda, D., Yan, L., Pettini, M., Shapley, A. E., Erb, D. K., & Adelberger, K. L. 2006, *ApJ*, 644, 792
- Rigby, J. R., et al. 2008, *ApJ*, 675, 262
- Sanders, D. B., Soifer, B. T., Elias, J. H., Madore, B. F., Matthews, K., Neugebauer, G., & Scoville, N. Z. 1988, *ApJ*, 325, 74
- Sanders, D. B., & Mirabel, I. F. 1996, *ARA&A*, 34, 749
- Swinbank, M., et al. 2010, *arXiv:1002.2518*
- Tacconi, L. J., et al. 2010, *Nature*, 463, 781
- Tran, Q. D., et al. 2001, *ApJ*, 552, 527
- van der Wel, A., Holden, B. P., Zirm, A. W., Franx, M., Rettura, A., Illingworth, G. D., & Ford, H. C. 2008, *ApJ*, 688, 48
- van Dokkum, P. G., et al. 2010, *ApJ*, 709, 1018
- van Dokkum, P. G., Kriek, M., Rodgers, B., Franx, M., & Puxley, P. 2005, *ApJ*, 622, L13
- Virani, S. N., Treister, E., Urry, C. M., & Gawiser, E. 2006, *AJ*, 131, 2373
- Wang, B., & Heckman, T. M. 1996, *ApJ*, 457, 645
- Weiß, A., et al. 2009, *ApJ*, 707, 1201
- Wuyts, S., Labbé, I., Schreiber, N. M. F., Franx, M., Rudnick, G., Brammer, G. B., & van Dokkum, P. G. 2008, *ApJ*, 682, 985
- Armus, L. 2009, *Astronomical Society of the Pacific Conference Series*, 408, 105
- Baldwin, J. A., Phillips, M. M., & Terlevich, R. 1981, *PASP*, 93, 5
- Bertin, E., & Arnouts, S. 1996, *A&AS*, 117, 393
- Bezanson, R., van Dokkum, P. G., Tal, T., Marchesini, D., Kriek, M., Franx, M., & Coppi, P. 2009, *ApJ*, 697, 1290
- Bruzual, G., & Charlot, S. 2003, *MNRAS*, 344, 1000
- Buat, V., et al. 2007, *ApJS*, 173, 404
- Buat, V., et al. 2005, *ApJ*, 619, L51
- Calzetti, D., Armus, L., Bohlin, R. C., Kinney, A. L., Koornneef, J., & Storch-Bergmann, T. 2000, *ApJ*, 533, 682
- Calzetti, D. 1997, *AJ*, 113, 162
- Caputi, K. I., et al. 2007, *ApJ*, 660, 97
- Chapman, S. C., Windhorst, R., Odewahn, S., Yan, H., & Conselice, C. 2003, *ApJ*, 599, 92
- Chary, R., & Elbaz, D. 2001, *ApJ*, 556, 562
- Daddi, E., et al. 2007, *ApJ*, 670, 173
- Daddi, E., et al. 2007, *ApJ*, 670, 156
- Dale, D. A., & Helou, G. 2002, *ApJ*, 576, 159
- Damen, M., Labbé, I., Franx, M., van Dokkum, P. G., Taylor, E. N., & Gawiser, E. J. 2009, *ApJ*, 690, 937
- Dasyra, K. M., Yan, L., Helou, G., Surace, J., Sajina, A., & Colbert, J. 2008, *ApJ*, 680, 232
- Dekel, A., et al. 2009, *Nature*, 457, 451
- Devlin, M. J., et al. 2009, *Nature*, 458, 737
- Donley, J. L., Rieke, G. H., Pérez-González, P. G., Rigby, J. R., & Alonso-Herrero, A. 2007, *ApJ*, 660, 167
- Dye, S., et al. 2009, *ApJ*, 703, 285
- Elbaz, D., Cesarsky, C. J., Chanial, P., Aussel, H., Franceschini, A., Fadda, D., & Chary, R. R. 2002, *A&A*, 384, 848
- Franx, M., van Dokkum, P. G., Schreiber, N. M. F., Wuyts, S., Labbé, I., & Toft, S. 2008, *ApJ*, 688, 770
- Genzel, R., et al. 2008, *ApJ*, 687, 59
- Genzel, R., et al. 1998, *ApJ*, 498, 579
- Hopkins, A. M., Connolly, A. J., Haarsma, D. B., & Cram, L. E. 2001, *AJ*, 122, 288
- Hopkins, P. F., Younger, J. D., Hayward, C. C., Narayanan, D., & Hernquist, L. 2010, *MNRAS*, 402, 1693
- Juneau, S., et al. 2005, *ApJ*, 619, L135
- Kartaltepe, J. S., et al. 2010, *ApJ*, 709, 572
- Kennicutt, R. C., et al. 2009, *ApJ*, 703, 1672
- Kennicutt, R. C., Jr. 1998, *ApJ*, 498, 541
- Kriek, M., van Dokkum, P. G., Franx, M., Illingworth, G. D., & Magee, D. K. 2009, *ApJ*, 705, L71
- Kriek, M., van Dokkum, P. G., Labbé, I., Franx, M., Illingworth, G. D., Marchesini, D., & Quadri, R. F. 2009, *ApJ*, 700, 221
- Kriek, M., et al. 2008, *ApJ*, 677, 219
- Kriek, M., et al. 2007, *ApJ*, 669, 776
- Le Floc'h, E., et al. 2005, *ApJ*, 632, 169
- Lonsdale, C. J., et al. 2003, *PASP*, 115, 897
- Luo, B., et al. 2008, *ApJS*, 179, 19

- Lutz, D., Valiante, E., Sturm, E., Genzel, R., Tacconi, L. J., Lehnert, M. D., Sternberg, A., & Baker, A. J. 2005, *ApJ*, 625, L83
- Lutz, D., Spoon, H. W. W., Rigopoulou, D., Moorwood, A. F. M., & Genzel, R. 1998, *ApJ*, 505, L103
- Magnelli, B., Elbaz, D., Chary, R. R., Dickinson, M., Le Borgne, D., Frayer, D. T., & Willmer, C. N. A. 2009, *A&A*, 496, 57
- Marcillac, D., Elbaz, D., Chary, R. R., Dickinson, M., Galliano, F., & Morrison, G. 2006, *A&A*, 451, 57
- Menéndez-Delmestre, K., et al. 2009, *ApJ*, 699, 667
- Murphy, E. J., Chary, R.-R., Alexander, D. M., Dickinson, M., Magnelli, B., Morrison, G., Pope, A., & Teplitz, H. I. 2009, *ApJ*, 698, 1380
- Muzzin, A., Marchesini, D., van Dokkum, P. G., Labbé, I., Kriek, M., & Franx, M. 2009, *ApJ*, 701, 1839
- Papovich, C., Rudnick, G., Rigby, J. R., Willmer, C. N. A., Smith, J.-D. T., Finkelstein, S. L., Egami, E., & Rieke, M. 2009, *ApJ*, 704, 1506
- Papovich, C., et al. 2007, *ApJ*, 668, 45
- Pérez-González, P. G., et al. 2005, *ApJ*, 630, 82
- Pope, A., et al. 2008, *ApJ*, 675, 1171
- Reddy, N. A., Erb, D. K., Pettini, M., Steidel, C. C., & Shapley, A. E. 2010, *arXiv:1002.0837*
- Reddy, N. A., Steidel, C. C., Fadda, D., Yan, L., Pettini, M., Shapley, A. E., Erb, D. K., & Adelberger, K. L. 2006, *ApJ*, 644, 792
- Rigby, J. R., et al. 2008, *ApJ*, 675, 262
- Sanders, D. B., Soifer, B. T., Elias, J. H., Madore, B. F., Matthews, K., Neugebauer, G., & Scoville, N. Z. 1988, *ApJ*, 325, 74
- Sanders, D. B., & Mirabel, I. F. 1996, *ARA&A*, 34, 749
- Swinbank, M., et al. 2010, *arXiv:1002.2518*
- Tacconi, L. J., et al. 2010, *Nature*, 463, 781
- Tran, Q. D., et al. 2001, *ApJ*, 552, 527
- van der Wel, A., Holden, B. P., Zirm, A. W., Franx, M., Rettura, A., Illingworth, G. D., & Ford, H. C. 2008, *ApJ*, 688, 48
- van Dokkum, P. G., et al. 2010, *ApJ*, 709, 1018
- van Dokkum, P. G., Kriek, M., Rodgers, B., Franx, M., & Puxley, P. 2005, *ApJ*, 622, L13
- Virani, S. N., Treister, E., Urry, C. M., & Gawiser, E. 2006, *AJ*, 131, 2373
- Wang, B., & Heckman, T. M. 1996, *ApJ*, 457, 645
- Weiβ, A., et al. 2009, *ApJ*, 707, 1201
- Wuyts, S., Labbé, I., Schreiber, N. M. F., Franx, M., Rudnick, G., Brammer, G. B., & van Dokkum, P. G. 2008, *ApJ*, 682, 985